



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

In (A), 42 is any one of the first set, and the remainder of the rows must be all different both within themselves and from each other, as written.

Proceeding to (B), after choosing say, 41, from the first set, each row must contain one and only one element from each row of (A). Since the elements in the several rows of (A) are permutable, it is allowable to take the columns in (A) for the remainders of the rows of (B). For convenience call the result of omitting 42 from (A) the *square*. Further discussion is confined to this square. As the square rotated about its principal diagonal appears in (B), any *new* six must be chosen from it as one would obtain a term from a determinant. But (A) and (B) admit interchanges of both columns and rows, without losing even this reciprocal relation, so that any new six may become the principal diagonal. Hence 36, 29, 22, 15, 8, 1, may be taken as any new six, and is to be coupled with one of the elements 0, 40, 39, 38, 37, to make the fourteenth set of seven. So far 36 has been used three times, hence it must appear in four more sets of six, and the same is true of 29, 22, 15, 8, 1, while none of these six elements may appear in the same set. Thus twenty-four sets of six are accounted for in addition to the above fourteen sets of seven. The remaining sets of six, five in number, will come from the square, with the diagonal omitted.

The square possesses two transformations which are of use in reducing the labor of computing sets of six. The first is a translation along the diagonal, with provision for the edges, and is

$$\begin{array}{cccc} (1, & 8, & 15, & 22, & 29, & 36) & (2, & 9, & 16, & 23, & 30, & 31) & (3, & 10, & 17, & 24, & 25, & 32) \\ (4, & 11, & 18, & 19, & 26, & 33) & (5, & 12, & 13, & 20, & 27, & 34) & (6, & 7, & 14, & 21, & 28, & 35) \end{array} \quad (C).$$

The second is the rotation about the diagonal. By a method to be shown, it is possible to find all the consistent sets of four sixes which contain 36. Now (C) leaves invariant the fourteen sets of seven already found, but gives by successive applications to the sets containing 36, those containing 29, then 22, etc. There are fifty-six sets of four sixes for 36, but six of them are invariant under the rotation, coupled with (42, 41), the fourteen sets of seven being also invariant. The remaining fifty sets are, by the rotation, reducible to twenty-five, so that thirty-one sets of four sixes for 36 are to be used. Of course there must be used with these, the entire fifty-six sets for 29, 22, etc.

In computing the sets for 36 the outline is this:

$$\begin{array}{cc} 36, 28, (19, 20, 21, 23) & 36, 28, 19, (14, 17) \\ 36, 27, (19, 20, 23) & 36, 27, 20, (13, 16, 17) \\ 36, 26, (19, 21, 23) & 36, 26, 21, (13, 16, 17) \\ 36, 25, (20, 21, 23) & 36, 25, 23, (14, 16) \end{array} \quad \begin{array}{c} (D), \\ \\ (E), \end{array}$$

in which the parentheses in (D) indicate those elements of the third line of the square which are available for the third element of the respective rows in (D). This gives eleven partial results from (D), of which the first is

(E). In this way the total of fifty-six sets of four sixes for 36 is obtained.

The remainder of the work begins with the comparison of such of the thirty-one sets for 36 with each of the fifty-six sets for 29. Although in upwards of forty cases one obtains consistent sets of eight sixes, all of these cases fail at the trial with the sets for 22, showing that the problem is impossible.

PROBLEMS FOR SOLUTION.

ALGEBRA.

283. Proposed by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Solve $w+x+y+z=4a$, $w^2+x^2+y^2+z^2=4a^2+4b^2$, $w^3+x^3+y^3+z^3=4a^3+12ab^2$, $w^4+x^4+y^4+z^4=4a^4+4b^4+4c^4+24a^2b^2$.

GEOMETRY.

316. Proposed by J. STEWART GIBSON, Department of Physics, Wadleigh High School, New York City.

Determine the locus of the vertices of parabolas described by particles thrown off from the circumference of a uniformly revolving wheel.

CALCULUS.

239. Proposed by L. H. MacDONALD, A. M., Ph. D., Sometime Tutor in the University of Cambridge, Jersey City, N. J.

Of all triangles inscribed in a circle, find that which has the greatest perimeter.

MECHANICS.

202. Proposed by W. J. GREENSTREET, M. A., Editor of The Mathematical Gazette, Stroud, England.

Three equal, uniform, similar rods AB , BC , CD , freely jointed at B and C , are hung from a point by two equal strings attached at A and D . Find the position of equilibrium.

MISCELLANEOUS.

171. Proposed by W. J. GREENSTREET, M. A., Editor of The Mathematical Gazette, Stroud, England.

If $\lim_{x \rightarrow a} \frac{\phi(x)}{\psi(x)} = \lambda$, show $\lim_{x \rightarrow a} \left[\frac{\lambda}{\phi(x)} - \frac{1}{\psi(x)} \right] = \frac{\lambda\psi''(a) - \phi''(a)}{2\phi'(a)\psi'(a)}$.

ERRATA.

Page 97, line 10. Vol. XIII, for $x=y=w=$ etc., read $x=x_1=x_2=$ etc.
 Page 98, line 1, for $G+D+U$ read $G+D+U+B$, B taken from table.
 Page 97, in table add .008 to each number from 34 to 43 inclusive.